



Naval Medical Research Institute

Bethesda, MD 20814-5055

MMRI 89-48

June 1989

AD-A216 240

PROSTAGLANDIN D₂ MODULATES HUMAN NEUTROPHIL INTRACELLULAR CALCIUM FLUX AND INHIBITS SUPEROXIDE RELEASE VIA ITS RING CARBONYL

C. O. SIMPKINS
D. L. MAZOROW
S. T. ALAILIMA
E. A. TATE
M. JOHNSON
K. SHARIF
W. SWEAT
D. B. MILLAR

Approved for public release;
distribution is unlimited

Naval Medical Research
and Development Command
Bethesda, Maryland 20814-5044

Department of the Navy
Naval Medical Command
Washington, D.C. 20372-5210

DTIC
ELECTE
DEC 27 1989
S E D

89 12 27 1 16

NOTICES

The opinions and assertions contained herein are the private ones of the writer and are not to be construed as official or reflecting the views of the naval service at large.

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Naval Medical Research Institute. Additional copies may be purchased from:

National Technical Information Service
5285 Port Royal Road
Springfield, Virginia 22161

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

Defense Technical Information Center
Cameron Station
Alexandria, Virginia 22304-6145

TECHNICAL REVIEW AND APPROVAL

NMRI 89-48

The experiments reported herein were conducted according to the principles set forth in the current edition of the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animal Resources, National Research Council.

This technical report has been reviewed by the NMRI scientific and public affairs staff and is approved for publication. It is releasable to the National Technical Information Service where it will be available to the general public, including foreign nations.

K. SORENSEN, CAPT, MC, USN

Commanding Officer
Naval Medical Research Institute

REPORT DOCUMENTATION PAGE


1a. REPORT SECURITY CLASSIFICATION UNCL			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 89-48			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research Institute		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
6c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20814-5055			7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, DC 20372-5120	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Medical Research & Development Command		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 10814-5044			10. SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO. 63706N	PROJECT NO. M0095.001
			TASK NO. 1005	WORK UNIT ACCESSION NO. DN977556
11. TITLE (Include Security Classification) Prostaglandin D ₂ Modulates Human Neutrophil Intracellular Calcium Flux and Inhibits Superoxide Release Via Its Ring Carbonyl				
12. PERSONAL AUTHOR(S) Cuthbert O. Simpkins, Denise L. Mazorow, Sione T. Alailima, Elin A. Tate, Mark Johnson, Khareem Sharif, William Sweat, and David B. Millar				
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM TO 1987	14. DATE OF REPORT (Year, Month, Day) June 1989	
15. PAGE COUNT 14				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Prostaglandin D ₂ , Prostaglandin F ₂ alpha	
FIELD	GROUP	SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We compared the effects of prostaglandin D ₂ (PGD ₂), prostaglandin F ₂ alpha (PGF ₂ alpha) and various ketones on O ₂ ⁻ release by human neutrophils, which had been stimulated by N-formyl methionyl leucyl phenylalanine (FMLP). Our data suggests that the ring carbonyl of PGD ₂ is essential to its inhibitory effect on O ₂ ⁻ release, but the carbonyl group as a ketone, alone, is not sufficient. Using the fluorescent Ca ²⁺ probe, Fura-2AM, we found that PGD ₂ decreased the late phase of FMLP invoked changes in intracellular free Ca ²⁺ ([Ca] _i), but that PGF ₂ alpha had no effect. Since cAMP altered FMLP stimulated [Ca] _i in a pattern similar to PGD ₂ , we suggest that PGD ₂ inhibits O ₂ ⁻ release by increasing intracellular cAMP which in turn causes an increase in the rate of decline in [Ca] _i . Furthermore, the ring carbonyl of PGD ₂ is crucial to its effect on O ₂ ⁻ as well as on [Ca] _i . <i>Keywords: Calcium, Cations, Superoxide S. (16)</i>				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Regina E. Hunt, Command Editor			22b. TELEPHONE (Include Area Code) (202) 295-0198	22c. OFFICE SYMBOL SD/RSD/NMRI

Prostaglandin D₂ (PGD₂) is the prostanoid which is most abundantly released by stimulated alevolar mast cells(1). Previously, we demonstrated that the release of O₂- by FMLP stimulated human neutrophils was inhibited by PGD₂. We found that PGD₂ was more potent than PGF₂ alpha, and suggested that the crucial determinant of potency was the ring carbonyl group on PGD₂. We also found that PGD₂ did not compete with FMLP at the FMLP receptor site (2).

TABLE OF CONTENTS

Acknowledgement.....	1
Introduction & Methods.....	2
Results.....	4
Figures	
1.....	5
2.....	6
3.....	10
Tables	
I.....	6
II	7
III.....	8
Discussion.....	11
References.....	12

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



ACKNOWLEDGEMENT

This work was supported by the Naval Medical Research and Development Command Work Unit No. M0095.001-1005. The opinions and assertions contained herein are the private ones of the authors and should not be construed as reflecting the views of the U. S. Navy, the naval service at large, or the Department of Defense.

INTRODUCTION

In this present work, we tested this hypothesis that only the carbonyl group as a ketone was necessary for inhibition of O_2^- . We also extended our understanding of the mechanism of O_2^- inhibition by PGD_2 through observing the effect of PGD_2 on FMLP stimulated $[Ca]_i$.

METHODS

Isolation of Neutrophils

Blood was obtained by venipuncture from healthy, medication-free volunteers. Preservative-free heparin (Gibco, Grand Island, NY) was used at a concentration of 12.5 units per ml of blood for anticoagulation. Erythrocytes were removed by gravity sedimentation through 2% dextran (Pharmacia, Upsale, Sweden) for 30 min. Lymphocyte Separation Medium (Litton Bionetics, Kensington, MD) was layered under the supernatant and centrifuged at $300 \times g$, room temperature for 10 min. Residual erythrocytes were removed by hypotonic lysis and the cells were centrifuged again at $150 \times g$. The neutrophil pellet was suspended in Hank's Balanced Salt Solution (HBSS, Gibco, Grand Island, NY) to the desired concentration. This procedure yielded 96% neutrophils by Wright's stain with a viability of at least 95% by trypan blue dye exclusion.

Assay of O_2^-

The superoxide dismutase-inhibitable reduction of cytochrome C assay of Babior et. al. was employed (3). Neutrophils were preincubated for 15 min. with various concentrations of test reagents with HBSS as control. After preincubation, 500 μ l of neutrophil suspension was added to the reaction mixture in 12 x 75 mm polystyrene tubes containing 1,500 μ l HBSS, test reagents (250 μ l) at various concentrations and 50 μ M cytochrome C (Sigma, St. Louis, MO). The final cell concentration was 1×10^6 /ml. After addition of

25 μ l of FMLP at a final concentration of 10^{-6} or 10^{-7} M, or HBSS as control, the reactants were mixed and a 12 ml aliquot was placed into 12 x 75 mm plastic tubes which contained 10 μ l superoxide dismutase (SOD: Boehringer Mannheim, W. Germany) at a final concentration of 50 μ g/ml. All tubes were incubated at 37°C for 10 min. At the end of the incubation period 15 μ l of DOS, also at a final concentration of 50 μ g/ml, was added to the other tubes to which SOD was not added initially. Centrifugation was at 150 x g and the absorbance of the supernatant at 550 nm was determined. O_2^- release was calculated as the difference in absorbance between the tubes which received SOD at the beginning, divided by an concentrations of PGD_2 (SIGMA, St. Louis, MO) was 95% or greater. Assays were performed in duplicate.

Measurement of Intracellular Calcium

5×10^6 cells were labelled with the membrane permeant probe Fura-2AM (Calbiochem) at 37°C for 30 min. at a probe concentration of 1×10^{-6} M. The cells were washed by centrifugation at 300 x g and resuspended in Hanks/HEPES. Fluorescent measurements were performed at 37°C in a Perkin Elmer Model 44B spectrophotofluoremeter. The cells were continually stirred during measurement. Alternative excitation wavelengths of 335 nm and 362 nm (slit 4 nm) were employed and emissions was measured at 510 nm (slit 8 nm). Calcium ion concentrations were calculated by the technique of Grynkiewicz et. al. (4) using the constants of Martell and Smith (5). FMLP control spectra were run prior to and after each experimental scan. Since our resting calcium levels determined with FURA-2AM were so much lower we investigated whether Quin-1AM gave us the same value for $[Ca]_i$ as did FURA-2AM. Where we had a resting

[Ca]_i with FURA-2AM of 16 - 20 nM, with QUIN-2AM we had resting levels of [Ca]_i from several individuals similar to those reported by others of 250-450 nM. Investigation with other cell lines gave similar results (manuscript in preparation, D.L. Mazorow and D.B. Millar).

Reagents

All reagents were prepared freshly each day. FMLP, prostaglandins, lipophilic ketones, and phorbol myristate acetate (PMA) were dissolved in dimethylsulfoxide (DMSO). Control solutions contained equal amounts of DMSO. Preliminary experiments showed no effect of DMSO on O₂- release (0.1 - 0.4%). The effect of PGD₂ upon the initial rate of reaction and the maximum O₂- released was measured as follows: The reaction mixture contained 3×10^6 cells/ml and 50 μ M cytochrome C in a total volume of 3 ml. Preincubation was in plastic cuvettes at 37°C for 15 min. with various concentrations of test reagents. The cuvettes were then placed into a temperature-controlled dual beam spectrophotometer with a magnetic stirrer. FMLP at 10^{-6} or 10^{-7} M or PMA at 4×10^{-7} M or greater was added to one cuvette and control solution added to the other. The reaction was monitored until a maximum was reached, which for FMLP took 5-8 min. and with PMA took as long as 60 min., when low concentrations (4×10^{-9} M) were used.

Statistical Analysis

All experiments were of a paired design. The paired Student's t-test employed. $P < 0.05$ was considered significant.

RESULTS

Previous experiments showed no effect of PGD₂ on the release of O₂- generated by a cell-free xanthine-xanthine oxidase system (2).

In six experiments performed with neutrophils from six individuals a

biphasic dose response curve of the inhibition of O_2^- release by PGD_2 was found (Fig. 1). The first phase occurred between 10^{-9} and $10^{-5}M$ and showed an inhibitory effect of up to 64%. In this concentration range PGD_2 had no effect on FMLP binding (2). In the second phase between 10^{-5} and $10^{-4}M$, O_2^- release was practically eliminated. This phase corresponds to the concentration at which PGD_2 decreases FMLP binding (2).




FIGURE 1.

Results are expressed as the percent of O_2^- release \pm S.E. in the absence of PGD_2 . Unstimulated O_2^- release is subtracted from all values. Each point is the average of six experiments performed with cells from six individuals.

$PGF_{2\alpha}$ is identical to PGD_2 except that $PGF_{2\alpha}$ has an hydroxyl group instead of a carbonyl group at the C11 position. The structures are shown in Fig. 2.




FIGURE 2.

The importance of the ring carbonyl was demonstrated by directly comparing the effect of PGD_2 to that of $\text{PGF}_{2\alpha}$ on O_2 - release (Table I).

TABLE I

Comparison of the Effect of 10^{-7}M PGD_2 and 10^{-7}M $\text{PGF}_{2\alpha}$ on O_2 - Release in Response to FMLP (10^{-7}M)

<u>Subject</u>	<u>PGD_2</u>	<u>$\text{PGF}_{2\alpha}$</u>	<u>Ratio $\text{PGD}_2/\text{PGF}_{2\alpha}$</u>
A	50.0	86.4	0.58
B	51.0	98.5	0.52
C	8.1	91.5	0.089
D	46.6	94.1	0.50
E	<u>39.8</u>	<u>69.8</u>	<u>0.57</u>
<u>MEAN</u> \pm S.E.	39.1 ± 7.3	$88. \pm 4.5$	0.46 ± 0.9

$P < 0.005$ for significant difference of PGD_2 from $\text{PGF}_{2\alpha}$.

These results suggest that a major determinant of the inhibition of O_2^- release by PGD_2 is the carbonyl group. Therefore, we studied the effects of various ketones and alcohols on FMLP stimulated O_2^- release (Table II).

TABLE II

Effect of Ketones and Alcohols on O_2^- Release

<u>Reagent ($10^{-6}M$)</u>	<u>% of Control + S.E.</u>
Cyclopentanone	98.4 ± 6.0
Cyclopentanol	88.5 ± 5.1
2,4 dimethylcyclopentanone	74.6 ± 7.8
Methanol	81.5 ± 4.0
Acetone	87.7 ± 6.8
PGD_2	54.5 ± 3.3
$PGF_{2\alpha}$	74.2 ± 5.3

$10^{-6}M$ FMLP was used. O_2^- release without alcohols or ketones was used as the control. Assays were performed on 2 to 4 individuals for each reagent. Included with each experiment was PGD_2 and $PGF_{2\alpha}$ at $10^{-6}M$ as controls.

None of the ketones or alcohols caused a marked decrease in O_2^- release. Therefore, the carbonyl group alone as a ketone is not sufficient to inhibit O_2^- release to a large degree. 2,4 dimethylcyclopentanone is more lipophilic than cyclopentanone and decreased O_2^- release to 74% of control compared with 98% of control for cyclopentanone. This result suggests that increased lipophilicity is factor in the inhibitory effect of a ketone.

It is also of interest that in most cases PGD₂ inhibited O₂⁻ release by approximately 50% regardless of whether PGD₂ and FMLP were used at 10⁻⁷M to 20⁻⁶M. For unknown reasons subject C of Table I was much more sensitive to PGD₂ than other individuals tested.

We found that PGD₂ decreased both the initial rate and the maximum quantity of O₂⁻ release (Table III). PGD₂ alone caused no change in O₂⁻ over the 15 min. preincubation period (data not shown).

TABLE III

Effect of PGD₂ (10⁻⁶M) on the Initial Rate and Emax of O₂⁻ Release Stimulated by 10⁻⁶M FMLP.

<u>Subject</u>	<u>Initial Rate</u>	<u>Emax</u>
A	85.1	46.1
B	74.5	79.5
C	73.6	56.2
D	77.9	48.6
E	66.7	50.3
F	<u>55.0</u>	<u>41.5</u>
Mean ± S.E.	72.1 ± 4.21	53.7 ± 5.5

Control initial rate was from 2.82 to 9.33 mMoles/Min/1.5 x 10⁷. Data are presented as percent of control cells. Control Emax was from 5.43 to 27.99 nMoles/1.5 x 10⁷ cells.

The reduction of the initial rate of O₂⁻ release by PGD₂ suggests that PGD₂ acted upon the rate limiting enzyme NADPH oxidase. This effect may be direct or indirect. In order to distinguish between these two possibilities, we determined the effect of PGD₂ on O₂⁻ release stimulated with PMA. PMA activates NADPH oxidase via a pathway which is different from that FMLP. PMA

directly activates protein kinase C, thereby bypassing the FMLP receptor and other early signal transduction mechanism (6). We reasoned that if PGD_2 were directly acting upon NADPH oxidase then its inhibitory effect on O_2^- release should be independent of the pathway of NADPH oxidase activation. We found that PGD_2 had no effect on PMA stimulated O_2^- release even at PMA concentrations as low as $4 \times 10^{-9}\text{M}$. Lower PMA concentrations did not stimulate O_2^- release. Therefore, PGD_2 is not acting directly upon NADPH oxidase (data not shown).

Next we examined the effect of PGD_2 on $[\text{Ca}]_i$ using the fluorescent dye Fura-2AM. PGD_2 had no effect on the initial rise in $[\text{Ca}]_i$, after stimulation by 10^{-6} FMLP. However, in the PDG_2 treated cells, the rate at which $[\text{Ca}]_i$ decreased toward baseline was more rapid than in control cells (Fig. 3A). Since prostaglandins have been shown to increase cAMP levels (7), we determined whether cAMP may mediate the changes in $[\text{Ca}]_i$ caused by PDG_2 . Incubation of cells with dibutyryl cAMP produced a change in rate of $[\text{Ca}]_i$ return to baseline similar to that of PGD_2 (Fig. 3). This experiment suggested that PGD_2 altered $[\text{Ca}]_i$ by altering cAMP levels. cAMP caused no change in the rate of return to baseline (data not shown).




FIGURE 3.

- A. Comparison of the effect of PGD_2 and $\text{PGF}_{2\alpha}$ on FMLP (10^{-6}M) induced changes in $[\text{Ca}]_i$. Resting $[\text{Ca}]_i$ levels are 16-20 nM. and FMLP induced peak $[\text{Ca}]_i$ levels are 80-110nM.
- B. The effect of 10^{-3} cAMP on changes in $[\text{Ca}]_i$ induced by 10^{-6} FMLP. All experiments were repeated three times with cells from at least three individuals and similar results were obtained.

DISCUSSION

Fanton and Kinnes (8) demonstrated that prostaglandins E_1 and I_2 inhibited FMLP stimulated O_2^- release in human neutrophils. They also showed that $PGF_{2\alpha}$ had no effect on O_2^- release (8). By comparing two prostaglandins which differed in only structural aspect, we showed that the carbonyl group of PGD_2 was essential to its inhibitory effect. The fact that a carbonyl as a ketone alone was not sufficient to inhibit O_2^- release was shown by the ineffectiveness of the ketones. Lipophilicity of the ketone may be important in determining potency, as demonstrated by the increased effect of 2,4 dimethylcyclopentanone over that of cyclopentanone.

Our previous experiments showed that concentrations equal to and less than $10^{-5}M$ PGD_2 had no effect on FMLP binding (2). Therefore, PGD_2 is not acting via the FMLP receptor. Togni et al. showed that 10^{-5} cAMP caused a more rapid recovery of $[Ca]i$ levels back to baseline but did not relate this observation to inhibition of O_2^- release. Our results suggest that PGD_2 may inhibit O_2^- release via cAMP. We found that cAMP but not cGMP causes the same changes in $[Ca]i$ as does PGD_2 . Furthermore, $PGF_{2\alpha}$ which had little effect on O_2^- release also had no effect on $[Ca]i$. The experiments with PMA, in which PGD_2 had no effect even at a PMA concentration as low as $4 \times 10^{-9}M$ showed that the PGD_2 interacts with the neutrophil in a carbonyl dependent manner to effect an increase in intracellular cAMP which enhances the recovery of kinase C and other kinases are Ca^{+2} dependent, it is possible that the decline in $[Ca]i$ caused by PGD_2 leads to the inhibition of O_2^- release (9). The lipophilicity rendered by the side chains of PGD_2 may allow the molecule to enter a critical

area of the cell membrane. The hydroxyl groups at C9 and C15 may be important in anchoring the prostaglandins into an important position by hydrogen bonding. Further work on studying the structure/function relationships among prostaglandins is clearly needed.

REFERENCES

1. C.C. Hardy, C. Robinson, A.E. Tattersfield, and S.J. Holgate. N. England Journal of Medicine, 311:209-212 (1984).
2. C.O. Simpkins, S.T. Alailima, E.A. Tate, and M. Johnson, J. Surg. Res. 41:645-652 (1986).
3. B.M. Babior, R.S. Kignes, J.T. Curnette. J. Clin. Invest. 52:741-744 (1973).
4. G. Grynkiewicz, M. Poenie, and R.Y. Tsien. J. Biol. Chem. 260:3440-3450 (1985).
5. A.E. Martell and R.M. Smith. Critical Stability Constants, 1, Plenum Press, NU (1974).
6. J.E. Niedel, L.J. Kuhn, G.R. Vanderbark. Proc. Nat'l Acad. Sci. USA 75:3822-3838 (1978).
7. T. Takenawaq, J. Ishitoya, Y. Nagai. J. Biol. Chem. 251:1092-1098 (1986).
8. J.C. Fantone and D.A. Kinnes. Biochem. Biophys. Res. Commun. 113:506-512 (1986).
9. P. Togni, G. Cabrini, F. Divirgilio. Biochem. J. 224:629-635 (1984).